



Measured correlation of nanoparticle magnetic moment and hydrodynamic size

Fock, Jeppe; Balceris, Christoph ; Costo, Rocio ; Zeng, Lunjie; Ludwig, Frank; Hansen, Mikkel Foug

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Fock, J., Balceris, C., Costo, R., Zeng, L., Ludwig, F., & Hansen, M. F. (2018). *Measured correlation of nanoparticle magnetic moment and hydrodynamic size*. Abstract from 12th International Conference on the Scientific and Clinical Applications of Magnetic Carriers, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Measured correlation of nanoparticle magnetic moment and hydrodynamic size

Jeppe Fock^{1*}, Christoph Balceris², Rocio Costo³, Lunjie Zeng⁴, Frank Ludwig², and Mikkel F. Hansen¹

¹ Department of Micro- and Nanotechnology, Technical University of Denmark, DK2800 Kongens Lyngby, Denmark

² Institute of Electrical Measurement and Fundamental Electrical Engineering, Technische Universität Braunschweig, Braunschweig D-38106, Germany

³ Instituto de Ciencia de Materiales de Madrid, ICMN/CSIC, Sor Juana Ines de la Cruz 3, 28049 Madrid, Spain

⁴ Department of Physics, Chalmers University of Technology, Gothenburg 41296, Sweden

* E-mail: jepf@nanotech.dtu.dk

The hydrodynamic size, V_h , and remanent magnetic moment, m , are of key importance for the properties of colloidal magnetic nanoparticle (MNP) dispersions. Often these parameters are treated independently although it is known that they are often correlated. In this study, we consider AC susceptibility (ACS) or optomagnetic (OM) measurements. ACS measures the magnetic response to an alternating magnetic field of frequency f and amplitude H , whereas the OM technique relies on measurements of the field-induced modulation of the optical transmission signal for MNPs with linked magnetic and optical anisotropies.¹ Using either of these techniques, the distributions of V_h or m can be estimated from low-field measurements vs. frequency or from low-frequency measurements vs. field,¹ but the correlation between V_h and m cannot be obtained.

Here, we present a method to determine the distributions of m and V_h as well as their correlation from dynamic magnetic measurements vs. frequency, f , and magnetic field strength, H . In this method the complete set of OM and/or ACS measurements are analysed in terms of a bivariate lognormal distribution of m and Brownian relaxation frequencies $f_B = k_B T / (6\pi\eta V_h)$, where η is the viscosity. The bivariate distribution accounts for the distributions of m and V_h and it assumes a power law relation between m and V_h .²

Fig. 1a shows ACS and OM measurements on Micromod BNF 80 MNPs vs. f and H . The lines are the fit to the above model with the bivariate distribution shown in Fig. 1b with the correlation $m \propto V_h^{0.5}$.² This square-root dependence is characteristic for multi-core particles.³ As a novel feature, we show that the use of higher harmonics in the analysis extends the measurement window and allows for the robust determination of values of f_B larger than the maximum value of f .




Fig 1(a) ACS and OM measurements (points) vs. H and f on BNF 80 particles. The solid lines are the fit obtained from simultaneous analysis of all data to a bivariate distribution of Brownian relaxation frequencies and magnetic moments. Colours from blue to red correspond to increasing field amplitudes. **(b)** Resulting bivariate distribution function. The inset shows a transmission electron microscopy (TEM) image of a particle. **(c)** Size histogram obtained from TEM. The line is the number-weighted size distribution from **(b)**. Figures were adapted from Fock *et al.*²

- (1) Fock, J.; Jonasson, C.; Johansson, C.; Hansen, M. F. *Phys. Chem. Chem. Phys.* **2017**, *19* (13), 8802–8814.
- (2) Fock, J.; Balceris, C.; Costo, R.; Zeng, L.; Ludwig, F.; Hansen, M. F. *Nanoscale* **2018**, *10* (4), 2052–2066.
- (3) Ahrentorp, F.; Astalan, A.; Blomgren, J.; Jonasson, C.; Wetterskog, E.; Svedlindh, P.; Lak, A.; Ludwig, F.; van IJzendoorn, L. J.; Westphal, F.; Grüttner, C.; Gehrke, N.; Gustafsson, S.; Olsson, E.; Johansson, C. *J. Magn. Magn. Mater.* **2015**, *380*, 221–226.